arXiv:math/0103136v2 [math.CV] 31 Oct 2001

Integrable systems and effectivisation
# of Riemann theorem about domains of the complex plane

S.M.Natanzon

1. Consider a closed analytic curve γ in the complex plane and denote by D_+ and D_- the interior and exterior domains with respect to the curve. The point $z = 0$ is assumed to be in D_+. Then according to Riemann theorem there exists a function $w(z) = \frac{1}{r}z + \sum\limits_{j=0}^{\infty} p_j z^{-j}$, mapping D_- to the exterior of the unit disk $\{w \in \mathbb{C} \mid |w| > 1\}$. It is follow from [1] that this function is described by formula $\log w = \log z - \partial_{t_0}(\frac{1}{2}\partial_{t_0} + \sum\limits_{k\geqslant 1} \frac{z^{-k}}{k}\partial_{t_k})v$, where $v = v(t_0, t_1, \bar{t}_1, t_2, \bar{t}_2, ...)$ is a function from infinite number of variables

$$t_0 = \frac{1}{\pi}\int\limits_{D_+} d^2z, \quad t_k = \frac{1}{\pi k}\int\limits_{D_-} z^{-k} d^2z.$$

Moreover, this function satisfies the dispersionless Hirota equation for $2D$ Toda lattice hierarchy.

$$(z - \xi)e^{D(z)D(\xi)v} = ze^{-\partial_{t_0}D(z)v} - \xi e^{-\partial_{t_0}D(\xi)v}. \tag{1}$$

$$(\bar{z} - \bar{\xi})e^{\bar{D}(\bar{z})\bar{D}(\bar{\xi})v} = \bar{z}e^{-\partial_{t_0}\bar{D}(\bar{z})v} - \bar{\xi}e^{-\partial_{t_0}\bar{D}(\bar{\xi})v}. \tag{2}$$

$$1 - e^{-D(z)\bar{D}(\bar{\xi})v} = \frac{1}{z\bar{\xi}}e^{\partial_{t_0}(\partial_{t_0} + D(z) + \bar{D}(\bar{\xi}))v}, \tag{3}$$

where

$$D(z) = \sum_{k\geqslant 1}\frac{z^{-k}}{k}\partial_{t_k}, \quad \bar{D}(\bar{z}) = \sum_{k\geqslant 1}\frac{\bar{z}^{-k}}{k}\partial_{\bar{t}_k}.$$

Thus for an effectivisation of Riemann theorem it is sufficiently to find a representation of v in the form of Taylor series

$$v = \sum N(i_0|i_1, ..., i_k|\bar{i}_1, ..., \bar{i}_{\bar{k}})t_0 t_{i_1}, ..., t_k \bar{t}_{\bar{i}_1}, ..., \bar{t}_{\bar{i}_{\bar{k}}}.$$

The numbers $N(i_0\big|i_1,...,i_k\big|\bar{i}_1,...,\bar{i}_{\bar{k}})$ for $i_\alpha, \bar{i}_\beta \leqslant 2$ is found in [1]. In this paper we find some recurrence relations, that give a possible to find all $N(i_0\big|i_1,...,i_k|\bar{i}_1,...,\bar{i}_{\bar{k}})$. For this we find some formulas for reconstruction any solutions of (1) – (3) via arbitrary Cauchy data $\frac{\partial^2 v}{\partial t_0 \partial t_i}\Big|_{t_0}, \frac{\partial^2 v}{\partial t_0 \partial \bar{t}_i}\Big|_{t_0}, \frac{\partial^2 v}{dt^2}\Big|_{t_0}$ (here $F\Big|_{t_0} = F\Big|_{t_1=\bar{t}_1=t_2=\bar{t}_2=\cdots=0}(t_0)$) and find the values of these data for our v.

2. Let us put $\partial_i = \partial_{t_i} = \frac{\partial}{\partial t_i}$ and $\bar{\partial}_i = \partial_{\bar{t}_i} = \frac{\partial}{\partial \bar{t}_i}$.

Lemma 1. $z - \sum\limits_{j=1}^{\infty} \frac{1}{j} z^{-j} \partial_1 \partial_j v = z e^{-\partial_0 D(z)} v$ *and*

$$\partial_1 \partial_j v = \sum_{m=1}^{\infty} \frac{(-1)^{m+1}}{m!} \sum_{k_1+\cdots+k_m=j+1} \frac{j}{k_1 \cdots k_m} \partial_0 \partial_{k_1} v \cdots \partial_0 \partial_{k_m} v.$$

Proof: According (1) $(z-\xi)e^{D(z)D(\xi)v} = (z-\xi)(1+(D(z)D(\xi)v)+\frac{1}{2}(D(z)D(\xi)v)^2+\cdots) = (z-\xi)(1+z^{-1}\xi^{-1}\partial_1^2 v + z^{-1}\sum\limits_{j=2}^{\infty}\frac{1}{j}\xi^{-j}\partial_1\partial_j v + \xi^{-1}\sum\limits_{j=2}^{\infty}\frac{1}{j}z^{-j}\partial_1\partial_j v + z^{-2}\xi^{-2}F) = (z-\xi) + \xi^{-1}\partial_1^2 v - z^{-1}\partial_1^2 v + \sum\limits_{j=2}^{\infty}\frac{1}{j}\xi^{-j}\partial_1\partial_j v - \sum\limits_{j=2}^{\infty}\frac{1}{j}z^{-j}\partial_1\partial_j v + z^{-1}\xi^{-1}F.$

On the other hand according (1) the function $(z-\xi)e^{D(z)D(\xi)v}$ is a sum of two functions $f_1(z) + f_2(\xi)$. Thus $F = 0$ and $ze^{-\partial_0 D(z)}v = z - \sum\limits_{j=1}^{\infty}\frac{1}{j}z^{-j}\partial_1\partial_j v$. Therefore,

$$\sum_{j=1}^{\infty}\frac{1}{j}z^{-(j+1)}\partial_1\partial_j v = 1 - e^{-\partial_0 D(z)}v = 1 - \big(1 + \sum_{m=1}^{\infty}\frac{(-\partial_0 D(z)v)^m}{m!}\big) =$$

$$-\sum_{m=1}^{\infty}\frac{(-1)^m}{m!}\Big(\sum_{k=1}^{\infty}\frac{z^{-k}}{k}\partial_0\partial_k v\Big)^m =$$

$$= -\sum_{m=1}^{\infty}\frac{(-1)^m}{m!}\Big(\sum_{n=1}^{\infty}z^{-n}\sum_{k_1+\cdots+k_m=n}\frac{1}{k_1\ldots k_m}\partial_0\partial_{k_1}v\cdots\partial_0\partial_{k_m}v\Big) =$$

$$= -\sum_{n=1}^{\infty}z^{-n}\Big(\sum_{m=1}^{\infty}\frac{(-1)^m}{m!}\sum_{k_1+\cdots+k_m=n}\frac{1}{k_1\ldots k_m}\partial_0\partial_{k_1}v\cdots\partial_0\partial_{k_m}v\Big).$$

Thus

$$\frac{1}{j}\partial_1\partial_j v = -\sum_{m=1}^{\infty}\frac{(-1)^m}{m!}\sum_{k_1+\cdots+k_m=j+1}\frac{1}{k_1\ldots k_m}\partial_0\partial_{k_1}v\cdots\partial_0\partial_{k_m}v.\square$$

Lemma 2. *$\partial_i\partial_j v = \sum\limits_{m=1}^{\infty}\sum\limits_{\substack{s_1+\cdots+s_m=j+1\\ s_i>1}}\frac{(-1)^{m+1}}{m}\frac{ij}{(s_1-1)\cdots(s_m-1)}P_{ij}(s_1-1,...,s_m-1)\cdot$*

$\cdot\partial_1\partial_{s_1-1}v\cdots\ \partial_1\partial_{s_m-1}v$, where $P_{ij}(s_1-1,...,s_m-1)$ is the number of representations of $\{i=i_1+\cdots+i_m|1\leqslant i_k\leqslant s_k-1,\ k=1,\cdots,m\}$ of number i.

Proof: According to lemma 1 and equation (1) $(z-\xi)e^{D(z)D(\xi)v}=$

$$=z-\sum_{j=1}^{\infty}\frac{1}{j}z^{-j}\partial_1\partial_j v-(\xi-\sum_{j=1}^{\infty}\frac{1}{j}\xi^{-j}\partial_1\partial_j v)=(z-\xi)-\sum_{j=1}^{\infty}\frac{1}{j}(z^{-j}-\xi^{-j})\partial_1\partial_j v.$$

Thus,

$$e^{D(z)D(\xi)}=1+z^{-1}\xi^{-1}\sum_{j=1}^{\infty}\frac{1}{j}\frac{(z^{-j}-\xi^{-j})}{(z^{-1}-\xi^{-1})}\partial_1\partial_j v=$$

$$=1+z^{-1}\xi^{-1}\sum_{j=1}^{\infty}\frac{1}{j}(\sum_{\substack{s+t=j-1\\ s,t\geqslant 0}}z^{-s}\xi^{-t})\partial_1\partial_j v=$$

$$=1+\sum_{j=1}^{\infty}\frac{1}{j}(\sum_{\substack{s+t=j+1\\ s,t\geqslant 1}}z^{-s}\xi^{-t})\partial_1\partial_j v.$$

Therefore,

$$D(z)D(\xi)v=\sum_{m=1}^{\infty}\frac{(-1)^{m+1}}{m}(\sum_{n=1}^{\infty}(\sum_{\substack{s+t=n+1\\ s,t\geqslant 1}}z^{-s}\xi^{-t})\frac{1}{n}\partial_1\partial_n v)^m=$$

$$=\sum_{j=1}^{\infty}\frac{(-1)^{m+1}}{m}\sum_{i,j\geqslant 1}z^{-i}\xi^{-j}.$$

$$\Big(\sum_{\substack{i_1+\cdots+i_m=i\\ j_1+\cdots j_m=j\\ i_k, j_k \geqslant 1}} \frac{1}{i_1+j_1-1}\partial_1\partial_{i_1+j_1-1}v\cdots\frac{1}{i_m+j_m-1}\partial_1\partial_{i_m+j_m-1}v\Big),$$

that is

$$\partial_i\partial_j v=\sum_{m=1}^{\infty}\sum_{s_1+\cdots+s_m=i+j}\frac{(-1)^{m+1}}{m}\frac{ij}{(s_1-1)\cdots(s_m-1)}\cdot$$

$$\cdot P_{ij}(s_1-1,...,s_m-1)\partial_1\partial_{s_1-1}v\cdots\partial_1\partial_{s_m-1}v.\Box$$

Remark. The equations

$$\partial_i\partial_j v=\sum_{m=1}^{\infty}\sum_{s_1+\cdots+s_m=i+j}\frac{(-1)^{m+1}}{m}\frac{ij}{(s_1-1)\cdots(s_m-1)}\cdot$$

$$\cdot P_{ij}(s_1-1,...,s_m-1)\partial_1\partial_{s_1-1}v\cdots\partial_1\partial_{s_m-1}v$$

describe the dispersionless limit of KP equation. Some other description of this hierarchy is presented in [2]. A comparison of these descriptions gives some nontrivial combinatorial identity P_{ij}.

Lemma 3. $\partial_i\partial_j v = \sum\limits_{m=1}^{\infty}\sum\limits_{p_1+\cdots+p_m=j+i}\frac{ij}{p_1\cdots p_m}T_{ij}(p_1\cdots p_m)\partial_0\partial_{p_1}v\cdots\partial_0\partial_{p_m}v,$ *where* $T_{ij}(p_1...p_m)=\sum\limits_{\substack{n_1+\cdots+n_k=m\\ n_i>0}}\frac{(-1)^{m+1}}{k}\frac{1}{n_1!\cdots n_k!}P_{ij}(p_1+\cdots+p_{q_1}-1,...,p_{q_{k-1}+1}+\cdots+p_{q_k}-1),$ *and* $q_j=\sum\limits_{i=1}^{j}n_i$.

Proof: According to lemmas 1 and 2

$$\partial_i\partial_j v=\sum_{m=1}^{\infty}\sum_{s_1+\cdots+s_m=j+i}\frac{(-1)^{m+1}}{m}\frac{ij}{(s_1-1)\cdots(s_m-1)}P_{ij}(s_1-1,\cdots,s_m-1)\cdot$$

$$\cdot\partial_1\partial_{s_1-1}v\cdots\partial_1\partial_{s_m-1}v)=\sum_{m=1}^{\infty}\sum_{s_1+\cdots+s_m=j+i}\frac{(-1)^{m+1}}{m}\frac{ij}{(s_1-1)\cdots(s_m-1)}\cdot$$

$$\cdot P_{ij}(s_1-1,\cdots,s_m-1)\Big(\sum_{n_1=1}^{\infty}\sum_{p_1+\cdots+p_{n_1}=s_1}\frac{(-1)^{n_1+1}}{n_1!}\frac{s_1-1}{p_1\cdots p_{n_1}}\partial_0\partial_{p_1}v\cdots\partial_0\partial_{p_{n_1}}v\Big)\cdots$$

$$\cdots\Big(\sum_{n_m=1}^{\infty}\sum_{p_1+\cdots+p_{n_m}=s_m}\frac{(-1)^{n_m+1}}{n_m!}\frac{s_m-1}{p_1\cdots p_{n_m}}\partial_0\partial_{p_1}v\cdots\partial_0\partial_{p_{n_m}}v\Big)=$$

$$=\sum_{m=1}^{\infty}\sum_{p_1+\cdots+p_m=j+i}\frac{ij}{p_1\cdots p_m}T_{ij}(p_1\cdots p_m)\partial_0\partial_{p_1}v\cdots\partial_0\partial_{p_m}v.\Box$$

Using the induction, we get from lemma 3

Lemma 4.

$$\partial_{i_1}\partial_{i_2}\cdots\partial_{i_k}v=\sum_{m=1}^{\infty}\Big(\sum_{\substack{s_1+\cdots+s_m=i_1+\cdots+i_k\\ \ell_1+\cdots+\ell_m=m+k-2\\ s_j,\ell_j\geqslant 1}}\frac{i_1\cdots i_k}{s_1\cdots s_m}\cdot$$

$$\cdot T_{i_1\cdots i_k}\begin{pmatrix}s_1\cdots s_m\\ \ell_1\cdots\ell_m\end{pmatrix}\partial_0^{\ell_1}\partial_{s_1}v\cdots\partial_0^{\ell_m}\partial_{s_m}v\Big),$$

$$\textit{where}\quad T_{i_1,i_2}\begin{pmatrix}s_1\cdots s_m\\ \ell_1\cdots\ell_m\end{pmatrix}=\begin{cases}T_{i_1i_2}(s_1\cdots s_m), & \textit{if } \ell_1=\cdots=\ell_m=1\\ 0 & \textit{in another cases}\end{cases},$$

$$T_{i_1\cdots i_k}\begin{pmatrix}s_1\cdots s_m\\ \ell_1\cdots\ell_m\end{pmatrix}=\sum_{\substack{1\leqslant i\leqslant j\leqslant m\\ s,\ell>0}}T_{i_1\cdots i_{k-1}}\begin{pmatrix}s_1\cdots s_{i-1}&\begin{pmatrix}s\\ \ell\end{pmatrix}&s_{j+1}\cdots s_m\\ \ell_1\cdots\ell_{i-1}&&\ell_{j+1}\cdots\ell_m\end{pmatrix}\cdot$$

$$\cdot T_{s,i_k}(s_i,s_{i+1},...,s_j)\frac{\ell!}{(\ell_i-1)!\cdots(\ell_j-1)!},$$

$$\textit{and}\quad s=s_i+s_{i+1}+\cdots+s_j-i_k,\quad \ell=(\ell_i-1)+\cdots+(\ell_j-1).\Box$$

3. Define now the Cauchy data for v.

Lemma 5.

$$\partial_0v|_{t_0}=-t_0+t_0\ln t_0\quad \partial_kv|_{t_0}=0\quad \textit{for}\quad k>0.$$

Proof: Put $\gamma = \{z \in \mathbb{C} | |z| = R\}$. The Schwarz function for γ is $S(z) = R^2 z^{-1}$. Thus according to [1 (26)]

$$t_0 = \frac{1}{\pi} \int\limits_{|z| \leqslant R} d^2 z = \frac{\pi R^2}{\pi} = R^2.$$

$$t_k = \frac{1}{2\pi i k} \int\limits_{|z|=R} z^{-k} R^2 z^{-1} dz = \frac{1}{2\pi i k} R^2 \oint\limits_{\gamma} z^{-(k+1)} dz = 0 \quad \text{for} \quad k > 0$$

Therefore, according to [1 (8), (26)]

$$\partial_0 v \Big|_{t_0} = \frac{2}{\pi} \int\limits_{|z| \leqslant R} \ln |z| d^2 z = \frac{2}{\pi} \int\limits_0^R dr \int\limits_0^{2\pi} d\varphi \cdot r \cdot \ln |r| =$$

$$= \frac{2 \cdot 2\pi}{2\pi} \int\limits_0^R \ln |r| dr^2 = 2(r^2 \ln r \Big|_0^R - \int\limits_0^R r dr) =$$

$$= r^2 \ln r^2 \Big|_0^R - 2 \frac{1}{2} r^2 \Big|_0^R = R^2 \ln R^2 - R^2 = -t_0 + t_0 \ln t_0.$$

$$\partial_k v \Big|_{t_0} = \frac{1}{2\pi i} \oint\limits_{|z|=R} z^k R^2 z^{-1} dz = \frac{R^2}{2\pi i} \oint\limits_{\gamma} z^{k-1} dz = 0 \quad \text{for} \quad k > 0. \square$$

Lemma 6.

$$\partial_i \bar{\partial}_j v|_{t_0} = \begin{cases} 0 & \textit{for } i \neq j, \\ i t_0^i & \textit{for } i = j. \end{cases}$$

Proof: It is follow from lemma 5 that $\partial_0 \partial_k v \Big|_{t_0} = 0$ for $k > 0$ and $\partial_0^2 v|_{t_0} = \ln t_0$. Moreover according to (3)

$$1 - e^{-D(z)\bar{D}(\bar{\xi})v} = z^{-1} \xi^{-1} e^{\partial_0 (\partial_0 + D(z) + \bar{D}(\bar{\xi}))v}.$$

Thus

$$e^{\partial_0 (\partial_0 + D(z) + \bar{D}(\bar{\xi}))v} \Big|_{t_0} = t_0$$

and

$$-D(z)\bar{D}(\bar{\xi})v\Big|_{t_0} = \ln(1 - z^{-1}\bar{\xi}^{-1}t_0) = -\sum_{k=1}^{\infty} kz^{-k}\xi^{-k}t_0^k.$$

Therefore $\partial_i\partial_j v\big|_{t_0} = 0$ for $i \neq j$ and $\partial_i^2 v\big|_{t_0} = it_0^i$ □.

Lemma 7.

$$\partial_i\bar{\partial}_{i_1}\cdots\bar{\partial}_{i_k}v\Big|_{t_0} = \bar{\partial}_i\partial_{i_1}\cdots\partial_{i_k}v\big|_{t_0} = \begin{cases} 0, & \textit{if} \quad i_1 + \cdots + i_k \neq i \\ i_1\cdots i_k\frac{i!}{(i-k+1)!}t_0^{i-k+1}, & \textit{if}\; i = i_1 + \cdots + i_k \end{cases}.$$

Proof: The differentials ∂ and $\bar{\partial}$ occur in (1) – (3)symmetrically. This gives the first equality. Moreover according to lemmas 4 and 5,

$$\partial_{i_1}\partial_{i_2}\cdots\partial_{i_k}v = \frac{i_1\cdots i_k}{i}T_{i_1\cdots i_k}\binom{i}{k-1}\partial_0^{k-1}\partial_i v +$$

$$+\sum_{m=2}^{\infty}\Big(\sum_{\substack{s_1+\cdots+s_m = i_1+\cdots+i_k \\ \ell_1+\cdots\ell_m = m+k-2 \\ s_j, t_j \geqslant 1}} \frac{i_1\cdots i_k}{s_1\cdots s_m}T_{i_1\cdots i_k}\begin{pmatrix} s_1\cdots s_m \\ \ell_1\cdots\ell_m\end{pmatrix}\cdot$$

$$\cdot\partial_0^{\ell_1}\partial_{s_1}v\cdots\partial_0^{\ell_m}\partial_{s_m}v\Big) =$$

$$= \frac{i_1\cdots i_k}{i}T_{i_1\cdots i_k}\binom{i}{k-1}\partial_0^{k-1}\partial_i v = \frac{i_1\cdots i_k}{i}\partial_0^{k-1}\partial_i v,$$

where $i = i_1 + \cdots + i_k$. This equality and lemma 6 give the second equality in the affirmation of lemma 7. □

Lemma 8.

$$\partial_{i_1}\cdots\partial_{i_k}\bar{\partial}_{\bar{i}_1}\cdots\bar{\partial}_{\bar{i}_{\bar{k}}}v\Big|_{t_0} = \sum_{i=1}^{\infty} N_i(i_1\cdots i_k\big|\bar{i}_1\cdots\bar{i}_{\bar{k}})t_0^{i-(k+\bar{k})+2}, \; \textit{where}\; N_i\big(i_1\cdots i_k\big|\bar{i}_1\cdots\bar{i}_{\bar{k}}\big) = 0$$

if $\sum_{j=1}^{k} i_j \neq i$ *or* $\sum_{j=1}^{\bar{k}} \bar{i}_j \neq i$. *In opposite case,* $N_i\big(i_1 \cdots i_{i_k} \big| \bar{i}_1, \cdots \bar{i}_{\bar{k}}\big) =$

$$= \sum_{m=1}^{\infty} \Big(\sum_{\substack{s_1 + \cdots + s_m = i_1 + \cdots + i_k \\ \ell_1 + \cdots \ell_m = m + k - 2 \\ s_j, \ell_j \geqslant 1}} i_1 \cdots i_k \bar{i}_k \cdots \bar{i}_{\bar{k}} \cdot$$

$$\cdot T_{i_1 \cdots i_k} \begin{pmatrix} s_1 \cdots s_m \\ \ell_1 \cdots \ell_m \end{pmatrix} S_{\bar{i}_1 \cdots \bar{i}_{\bar{k}}} \Big(s_1, ..., s_m \Big) \Big),$$

where

$$S_{\bar{i}_1 \cdots \bar{i}_{\bar{k}}}(s_1, ..., s_m) = \sum_{n_1 + \cdots + n_m = k} \sum \frac{(s_1 - 1)! \cdots (s_m - 1)!}{(s_1 - n_1 + 1 - \ell_1)! \cdots (s_m - n_m + 1 - \ell_m)!}$$

and second sum is the sum by all partitions of the set $\{\bar{i}_1, ..., \bar{i}_{\bar{k}}\}$ *on subsets* $\{j_1^p, ..., j_{n_p}^p\}$ $(p = 1, ..., m)$ *such that* $\sum_{\alpha=1}^{n_p} j_\alpha^p = s_p$. $\Box$

Proof: According to lemma 4

$$\partial_{i_1} \cdots \partial_{i_k} \bar{\partial}_{\bar{i}_1} \cdots \bar{\partial}_{\bar{i}_{\bar{k}}} v = \bar{\partial}_{\bar{i}_{\bar{i}}} \cdots \bar{\partial}_{\bar{i}_{\bar{k}}} \Big(\sum_{m=1}^{\infty} \Big(\sum_{\substack{s_1 + \cdots + s_m = i_1 + \cdots + i_k \\ \ell_1 + \cdots + \ell_m = m + k - 2 \\ s_j, \ell_j \geqslant 1}} \frac{i_1 \cdots i_k}{s_1 \cdots s_m} \cdot$$

$$\cdot T_{i_1 \cdots i_k} \begin{pmatrix} s_1 \cdots s_m \\ \ell_1 \cdots \ell_m \end{pmatrix} \partial_0^{\ell_1} \partial_{s_1} v \cdots \partial_0^{\ell_m} \partial_{s_m} v \Big) \Big) =$$

$$= \sum \Big(\sum_{m=1}^{\infty} \Big(\sum_{\substack{s_1 + \cdots + s_m = i_1 + \cdots + i_k \\ \ell_1 + \cdots + \ell_m = m + k - 2 \\ s_j, \ell_j \geqslant 1}} \frac{i_1 \cdots i_k}{s_1 \cdots s_m} T_{i_1 \cdots i_k} \cdot$$

$$\cdot \begin{pmatrix} s_1 \cdots s_m \\ \ell_1 \cdots \ell_m \end{pmatrix} \partial_0^{\ell_1} \partial_{s_1} \bar{\partial}_{\bar{j}_1}^1 \cdots \bar{\partial}_{\bar{j}_{n_1}}^1 v \cdots \partial_0^{\ell_m} \partial_{s_m} \bar{\partial}_{\bar{j}_1}^m \cdots \bar{\partial}_{\bar{j}_{n_m}}^m v \Big) \Big),$$

where the first sum is taken by all partitions of $\{\bar{i}_1, ..., \bar{i}_{\bar{k}}\}$ on subset $\{(j_i^1, ..., j_{n_1}^1) ... (j_1^m, ..., j_{n_m}^m)\}$. According to lemma 7 this gives the affirmation of lemma 8 $\Box$.

Lemma 8 gives

Theorem.

$$v=\frac{1}{2}t_0^2\log t_0-\frac{3}{4}t_0+\sum_{\substack{i_1<\cdots<i_k\\ \bar{i}_1<\cdots\bar{i}_{\bar{k}}}}\sum_{n_j,\bar{n}_j=1}^{\infty}\frac{1}{n_1!\cdots n_k!\bar{n}_1!\cdots\bar{n}_{\bar{k}}!}N_i\left(\begin{matrix}i_1...i_k\\ n_1...n_k\end{matrix}\,\middle|\,\begin{matrix}\bar{i}_1...\bar{i}_{\bar{k}}\\ \bar{n}_1...\bar{n}_{\bar{k}}\end{matrix}\right)\cdot$$

$$\cdot t_0^{i-(\sum\limits_{i=1}^{k}n_i+\sum\limits_{i=1}^{\bar{k}}\bar{n}_i)+2}\quad t_{i_1}^{n_1}\cdots t_{i_k}^{n_k}\bar{t}_{\bar{i}_1}^{\bar{n}_1}\cdots\bar{t}_{\bar{i}_{\bar{k}}}^{\bar{n}_{\bar{k}}},$$

where

$$N_i\left(\begin{matrix}i_1...i_k\\ n_1...n_k\end{matrix}\,\middle|\,\begin{matrix}\bar{i}_1...\bar{i}_{\bar{k}}\\ \bar{n}_1...\bar{n}_{\bar{k}}\end{matrix}\right)=0,$$

if $i\neq\sum\limits_{j=1}^{k}n_ji_j$ *or* $i\neq\sum\limits_{j=1}^{\bar{k}}\bar{n}_j\bar{i}_j$. *In opposite case*

$$N_i\left(\begin{matrix}i_1...i_k\\ n_1...n_k\end{matrix}\,\middle|\,\begin{matrix}\bar{i}_1...\bar{i}_{\bar{k}}\\ |n_1...\bar{n}_{\bar{k}}\end{matrix}\right)=N_i(i_1...i_1i_2...i_2...i_k...i_k|\bar{i}_1...\bar{i}_1\bar{i}_2...\bar{i}_2...\bar{i}_{\bar{k}}...\bar{i}_{\bar{k}}),$$

where in the last parentheses any i_j *(respectively any* $\bar{i}_j$ *) occurs* n_j *(respectively* $\bar{n}_j$*) cases.*

Reference.

1. I.K.Kostov, I.Krichever, M.Mineev–Weinstein, P.B.Wiegmann, A.Zabrodin, τ-function for anatytic curves, arXiv:hep-th/0005259.

2. S.M.Natanzon, Formulas for A_n and B_n–solutions of WDVV equations, Journal of Geometry and Physics, (2001).